

**TECHNICAL DATA SHEET**
**Catalog # MSP43**
**PDH Kinase 3 (PDK3)**

Rev.0

**LOT #:**
**COMPONENTS:** 50 µg human recombinant PDK3

**APPLICATIONS:** Phosphorylation (inactivation) of PDH

**CONCENTRATION:** 0.5 mg/mL in 10 mM NaPO<sub>4</sub>, pH 7.5, 250 mM NaCl, 30% glycerol, 5 mM DTT, 0.5 mM EDTA, 0.05% Triton X-100


**STORAGE CONDITIONS:** Store at -80°C.

**COUNTRY OF ORIGIN:** USA

**BACKGROUND:**

Shifts in metabolism are often regulated by reversible phosphorylation of particular amino-acids within regulatory enzymes. The pyruvate dehydrogenase complex (PDH, or PDC) (E.C.1.2.4.1) is a key regulatory enzyme in carbohydrate metabolism. This multimeric enzyme complex is composed of five proteins: E1 $\alpha$ , E1 $\beta$ , E2, E3 and the E3 binding protein, which are present in a conserved stoichiometric manner. Tissue- and diet-specific regulation of PDH occur via four PDH kinases (E.C.2.7.11.2), PDK1, 2, 3 and 4, and two PDH phosphatases (E.C.3.1.3.43), PDP1 and 2.

The PDH kinases are members of the serine/threonine class of kinase enzymes. These ATP-dependent enzymes are bound to the E2 domain of PDH and phosphorylate three particular serine residues within the E1 $\alpha$  subunit, thus inhibiting enzyme activity (see Figure 1). Aberrant regulation of PDKs has been observed in metabolic diseases such as diabetes, and is also related to changes in cancer tumor genesis. As a result, there is much interest in identifying drug candidates that regulate PDH kinase activity.



**Figure 1.** Schematic of the PDH kinase reaction. ATP-dependent phosphorylation of PDH occurs at serines 232, 293, and 300 of the E1 $\alpha$  subunit. Phosphorylation results in inactivation of the PDH complex.

This product provides the researcher with active, human recombinant PDK3. His-tagged PDK3 was bacterial-expressed and column-purified. The enzyme is supplied in a buffer perfectly suited for use with our PDH enzyme activity assays (MSP18 and MSP30). Also available are immunocapture assays for measuring total PDH levels, and companion phospho-specific detector antibodies against all three phosphorylation sites. Please see [http://www.mitosciences.com/pyruvate\\_dehydrogenase.html](http://www.mitosciences.com/pyruvate_dehydrogenase.html) for a complete list of available products. For a review of PDH, including an overview of protocols for using the PDK's and PDP's with our PDH enzyme activity and protein quantity assays, please download our PDH Playbook ([http://www.mitosciences.com/pdh\\_playbook.pdf](http://www.mitosciences.com/pdh_playbook.pdf)).

**Note: This product is for research purposes only. It is not to be used in humans or for diagnostic purposes.**

WARRANTY – MitoSciences Inc warrants that products will perform as indicated in the published Technical Data Sheet for 6 months from date of purchase when stored according to specifications and when used consistent with recommended protocols. If you experience results which materially differ from those described, send evidence of the non-performing product for replacement of the original product purchased or a credit toward any other of the company's products or services.